

## FINAL NEET(UG)-2024 (EXAMINATION)

(Held On Sunday 5th MAY, 2024)

## **CHEMISTRY**

# Chemistry: Section-A (Q. No. 51 to 85)


- **51.** 'Spin only' magnetic moment is same for which of the following ions?
  - A. Ti<sup>3+</sup>
- B. Cr<sup>2+</sup>
- C. Mn<sup>2+</sup>
- D. Fe<sup>2+</sup>
- E. Sc<sup>3+</sup>

Choose the most appropriate answer from the options given below:

- (1) B and D only
- (2) A and E only
- (3) B and C only
- (4) A and D only

## Ans. (1)

**52.** The most stable carbocation among the following is:





Ans. (4)

## **TEST PAPER WITH ANSWER**

**53.** Given below are two statements:

**Statement-I**: The boiling point of hydrides of Group-16 elements follow the order

 $H_2O > H_2Te > H_2Se > H_2S$ .

**Statement-II**: On the basis of molecular mass,  $H_2O$  is expected to have lower boiling point than the other members of the group but due to the presence of extensive H-bonding in  $H_2O$ , it has higher boiling point.

In the light of the above statements, choose the *correct* answer from the options given below:

- (1) Both statement-I and Statement-II are true.
- (2) Both statement-I and Statement-II are false.
- (3) Statement-I is the true but Statement-II is false.
- (4) Statement-I is false but Statement-II is true.

#### Ans. (1)

**54.** Match List I with List II.

## List I

#### List II

(Compound)	(Shape/geometry)
(A) NH <sub>3</sub>	(I) Trigonal Pyramidal
(B) BrF <sub>5</sub>	(II) Square Planar
(C) XeF <sub>4</sub>	(III) Octahedral
(D) SF <sub>6</sub>	(IV) Square Pyramidal

Choose the correct answer from the options given below:

- (1) A-I, B-IV, C-II, D-II (2) A-II, B-IV, C-III, D-I
- (3) A-III, B-IV, C-I, D-II (4) A-II, B-III, C-IV, D-I

#### Ans. (1)

- **55.** The highest number of helium atoms is in:
  - (1) 4 mol of helium
  - (2) 4 u of helium
  - (3) 4 g of helium
  - (4) 2.271098 L of helium at STP



Identify the correct reagents that would bring about **56.** the following transformation

- (1) (i)  $H_2O/H^+$  (ii)  $CrO_3$
- (2) (i)  $BH_3$  (ii)  $H_2O_2 / OH$  (iii) PCC
- (3) (i) BH<sub>3</sub> (ii) H<sub>2</sub>O<sub>2</sub> /  $\overset{\circ}{O}$ H (iii) Alk. KMnO<sub>4</sub> (iv) H<sub>3</sub>O<sup> $\oplus$ </sup>
- (4) (i) H<sub>2</sub>O/H<sup>+</sup> (ii) PCC

Ans. (2)

**57**.

Match List I with L	ist II.	
List-I		List-II
(Process)		(Conditions)
A. Isothermal	I.	No heat exchange
process		
B. Isochoric	II.	Carried out at constant
process		temperature
C. Isobaric proc	ess III.	Carried out at constant
		volume
D. Adiabatic	IV.	Carried out at constant
process		pressure
Choose the correct	ct answe	r from the options given
below:		
(1) A-IV, B-III, C-II	, D-I	
(2) A-IV, B-II, C-III	, D-I	

- (3) A-I, B-II, C-III, D-IV
- (4) A-II, B-III, C-IV, D-I

Ans. (4)

**58.** Which one one of the following alcohols reacts instantaneously with Lucas reagent?

(4) 
$$CH_3 - C - OH$$
  
 $CH_3 - C - OH$   
 $CH_3$ 

Ans. (4)

In which of the following equilibria,  $K_p$  and  $K_c$  are **59**. **NOT** equal?

- (1)  $PCl_{5(g)} \rightleftharpoons PCl_{3(g)} + Cl_{2(g)}$
- (2)  $H_{2(q)} + I_{2(q)} \rightleftharpoons 2HI_{(q)}$
- (3)  $CO_{(g)} + H_2O_{(g)} \rightleftharpoons CO_{2(g)} + H_{2(g)}$
- $(4) \ 2BrCl_{(g)} \Longrightarrow Br_{2(g)} + Cl_{2(g)}$

Ans. (1)

**60**. Match List I with List II.

List I	List II
<b>Quantum Number</b>	Information provided
A. $m_{\ell}$	I. shape of orbital
B. $m_s$	II. size of orbital
C. <i>l</i>	III. orientation of orbital
D. n	IV. orientation of spin of
	electron

Choose the correct answer from the options given below:

- (1) A-I, B-III, C-II, D-IV
- (2) A-III, B-IV, C-I, D-II
- (3) A-III, B-IV, C-II, D-I
- (4) A-II, B-I, C-IV, D-III

Ans. (2)

**61**. Given below are two statements:

> Statement I: Aniline does not undergo Friedel-Crafts alkylation reaction

> **Statement II**: Aniline cannot be prepared through Gabriel synthesis.

> In the light of the above statements, choose the correct answer from the options given below:

- (1) Both Statement I and Statement II are true.
- (2) Both Statement I and Statement II are false.
- (3) Statement I is correct but Statement II is false.
- (4) Statement I is incorrect but Statement II is true.





Intramolecular hydrogen bonding is present in:

(4) HF

## Ans. (1)

- **63.** On heating, some solid substances change from solid to vapour state without passing through liquid state. The technique used for the purification of such solid substances based on the above principle is known as:
  - (1) Crystallization
  - (2) Sublimation
  - (3) Distillation
  - (4) Chromatography

## Ans. (2)

- 64. In which of the following processes entropy increases?
  - A. A liquid evaporates to vapour
  - B. Temperature of a crystalline solid lowered from 130 K to 0K.
  - C.  $2NaHCO_{3(s)} \rightarrow Na_2CO_{3(s)} + CO_{2(g)} + H_2O_{(g)}$
  - D.  $Cl_{2(q)} \rightarrow 2 Cl_{(q)}$

Choose the correct answer from the options given below:

- (1) A and C
- (2) A, B and D
- (3) A, C and D
- (4) C and D

#### Ans. (3)

- Among Group 16 elements, which one does **NOT** show -2 oxidation state?
  - (1) O

(2) Se

- (3) Te
- (4) Po

## Ans. (4)

Match List-II with List-II. 66.

## List-I (Conversion)

## List-II (Number of

## Faraday required)

- (A) 1 mol of  $H_2O$  to  $O_2$
- (B) 1 mol of  $MnO_4^-$  to  $Mn^{2+}$
- (C) 1.5 mole of Ca from molten CaCl<sub>2</sub>
- (II) 2F (III) 1F

(IV) 5F

(I) 3F

- (D) 1 mol of FeO to  $Fe_2O_3$ 
  - (1) A-II, B-IV, C-I, D-III
  - (2) A-III, B-IV, C-I, D-II
  - (3) A-II, B-III, C-I, D-IV
  - (4) A-III, B-IV, C-II, D-I

#### Ans. (1)

- **67**. Arrange the following elements in increasing order of electronegativity.
  - N, O, F, C, Si

Choose the correct answer from the options given

- (1) Si < C < N < O < F (2) Si < C < O < N < F
- (3) O < F < N < C < Si (4) F < O < N < C < Si

#### Ans. (1)

- **68.** A compound with a molecular formula of  $C_6H_{14}$  has two tertiary carbons. Its IUPAC name is:
  - (1) n-hexane
- (2) 2-methylpentane
- (3) 2, 3-dimethylbutane (4) 2, 2-dimethylbutane

#### Ans. (3)

- **69**. Fehling's solution 'A' is
  - (1) aqueous copper sulphate
  - (2) alkaline copper sulphate
  - (3) alkaline solution of sodium potassium tartrate (Rochelle's salt)
  - (4) aqueous sodium citrate

#### Ans. (1)


- **70**. Activation energy of any chemical reaction can be calculated if one knows the value of
  - (1) rate constant at standard temperature.
  - (2) probability of collision.
  - (3) orientation of reactant molecules during collision.
  - (4) rate constant at two different temperatures.


#### Ans. (4)





Which plot of  $\ln k$  vs  $\frac{1}{T}$  is consistent with

Arrhenius equation?










Ans. (4)


Match List I with List II. **72**.

> List I (Reaction)

List II (Reagents/Condition)

(A) 
$$\longrightarrow$$
 2  $\longrightarrow$  0 (I)  $\bigcirc$  Cl/Anhyd.AlCl<sub>3</sub>





(III) KMnO<sub>4</sub>/KOH,  $\Delta$ 

$$(D) \bigcirc CH_2CH_3 \bigcirc COOK$$

(IV) (i) O<sub>3</sub> (ii) Zn-H<sub>2</sub>O

Choose the correct answer from the options given below:

(1) A-IV, B-I, C-III, D-II

(2) A-III, B-I, C-II, D-IV

(3) A-IV, B-I, C-II, D-III

(4) A-I, B-IV, C-II, D-III

Ans. (3)

The compound that will undergo  $S_N^1$  reaction with **73.** the fastest rate is:

$$(2)$$
  $\longrightarrow$   $B_1$ 

Ans. (4)

- **74**. Which reaction is **NOT** a redox reaction?
  - (1)  $Zn + CuSO_4 \rightarrow ZnSO_4 + Cu$
  - (2)  $2 \text{ KClO}_3 + I_2 \rightarrow 2 \text{ KIO}_3 + \text{Cl}_2$
  - (3)  $H_2 + Cl_2 \rightarrow 2 HCl$
  - (4)  $BaCl_2 + Na_2SO_4 \rightarrow BaSO_4 + 2NaCl$

Ans. (4)

**75.** Given below are two statements:

> **Statement I:** The boiling point of three isomeric pentanes follows the order

n-pentane > isopentane > neopentane

**Statement II:** When branching increases, the molecule attains a shape of sphere. This results in smaller surface area for contact, due to which the intermolecular forces between the spherical molecules are weak, thereby lowering the boiling point.

In the light of the above statements, choose the most appropriate answer from the options given below:

- (1) Both statement I and Statement II are correct
- (2) Both Statement I and Statement II are incorrect
- (3) Statement I is correct but Statement II is incorrect
- (4) Statement I is incorrect but Statement II is correct

Ans. (1)

**76**. Given below are two statements:

> **Statement I**: Both  $[Co(NH_3)_6]^{+3}$  and  $[CoF_6]^{3-}$ complexes are octahedral but differ in their magnetic behaviour.

> **Statement II**: [Co(NH<sub>3</sub>)<sub>6</sub>]<sup>3+</sup> is diamagnetic whereas  $[CoF_6]^{3-}$  is paramagnetic.

In the light of the above statements, choose the correct answer from the options given below:

- (1) Both statement I and Statement II are true
- (2) Both Statement I and Statement II are false
- (3) Statement I is true but Statement II is false
- (4) Statement I is false but Statement II is true





#### **77.** Match List I with List II.

List-I	
Molecule	)

## List-II (Number and types of bond/s between two carbon atoms)

- A. ethane
- I. one  $\sigma$ -bond and two  $\pi$ -bonds
- B. ethene
- II. two  $\pi$ -bonds
- C. carbon molecule, C<sub>2</sub> III.
  - III. one  $\sigma$ -bond IV. one  $\sigma$ -bond and
- D. ethyne
- IV. one  $\sigma$ -bond and one  $\pi$ -bond

Choose the correct answer from the options given below:

- (1) A-I, B-IV, C-II, D-III
- (2) A-IV, B-III, C-II, D-I
- (3) A-III, B-IV, C-II, D-I
- (4) A-III, B-IV, C-I, D-II

## Ans. (3)

- **78.** The Henry's law constant ( $K_H$ ) values of three gases (A, B, C) in water are 145,  $2 \times 10^{-5}$  and 35 kbar, respectively. The solubility of these gases in water follow the order:
  - (1) B > A > C
- (2) B > C > A
- (3) A > C > B
- (4) A > B > C

#### Ans. (2)

- **79.** The energy of an electron in the ground state (n = 1) for  $He^+$  ion is -xJ, then that for an electron in n = 2 state for  $Be^{3+}$  ion in J is :
  - (1) x
- (2)  $-\frac{x}{9}$
- (3) 4x
- $(4) \frac{4}{9}x$

#### Ans. (1)

- **80.** The E° value for the  $Mn^{3+}/Mn^{2+}$  couple is more positive than that of  $Cr^{3+}/Cr^{2+}$  or  $Fe^{3+}/Fe^{2+}$  due to change of
  - (1) d<sup>5</sup> to d<sup>4</sup> configuration
  - (2) d<sup>5</sup> to d<sup>2</sup> configuration
  - (3) d<sup>4</sup> to d<sup>5</sup> configuration
  - (4) d<sup>3</sup> to d<sup>5</sup> configuration
- Ans. (3)

- **81.** The reagents with which glucose does **not** react to give the corresponding tests/products are
  - A. Tollen's reagent
- B. Schiff's reagent
- C. HCN
- D. NH<sub>2</sub>OH
- E. NaHSO<sub>3</sub>

Choose the correct options from the given below:

- (1) B and C
- (2) A and D
- (3) B and E
- (4) E and D

- Ans. (3)
- **82.** Match List I with List II.

# List-I (Complex) (Type of isomerism)

- A.  $\left[\text{Co}\left(\text{NH}_3\right)_5\left(\text{NO}_2\right)\right]\text{Cl}_2$
- I. Solvate isomerism
- B.  $\left[\text{Co(NH}_3)_5\left(\text{SO}_4\right)\right]$ Br
- II. Linkage isomerism
- $C. \ \left[ \text{Co(NH}_3)_6 \right] \! \left[ \text{Cr(CN)}_6 \right]$
- III. Ionization isomerism
- D.  $\left[\text{Co}\left(\text{H}_2\text{O}\right)_6\right]\text{Cl}_3$
- IV. Coordination isomerism

Choose the correct answer from the options given below:

- (1) A-II, B-III, C-IV, D-I
- (2) A-I, B-III, C-IV, D-II
- (3) A-I, B-IV, C-III, D-II
- (4) A-II, B-IV, C-III, D-I

#### Ans. (1)

- **83.** Arrange the following elements in increasing order of first ionization enthalpy:
  - Li, Be, B, C, N

Choose the correct answer from the options given below:

- (1) Li < Be < B < C < N
- (2) Li < B < Be < C < N
- (3) Li < Be < C < B < N
- (4) Li < Be < N < B < C
- Ans. (2)





- 1 gram of sodium hydroxide was treated with 84. 25 mL of 0.75 M HCL solution, the mass of sodium hydroxide left unreacted is equal to
  - (1)750 mg
- (2) 250 mg
- (3) Zero mg
- (4) 200 mg

Ans. (2)

**85.** For the reaction  $2A \Longrightarrow B + C$ ,  $K_c = 4 \times 10^{-3}$ . At a given time, the composition of reaction mixture is:  $[A] = [B] = [C] = 2 \times 10^{-3} M.$ 

Then, which of the following is correct?

- (1) Reaction is at equilibrium.
- (2) Reaction has a tendency to go in forward direction.
- (3) Reaction has a tendency to go in backward direction
- (4) Reaction has gone to completion in forward direction.

Ans. (3)

## Chemistry: Section-B (Q. No. 86 to 100)

- 86. Given below are certain cations. Using inorganic qualitative analysis, arrange them in increasing group number from 0 to VI.
  - A. Al<sup>3+</sup>
- B. Cu<sup>2+</sup>
- C. Ba<sup>2+</sup>
- D. Co<sup>2+</sup>
- E. Mg<sup>2+</sup>

Choose the correct answer from the options given below:

- (1) B, A, D, C, E
- (2) B, C, A, D, E
- (3) E, C, D, B, A
- (4) E, A, B, C, D

Ans. (1)

**87**. The products A and B obtained in the following reactions, respectively, are

$$3ROH + PCl_3 \rightarrow 3RCl + A$$

$$ROH + PCl_5 \rightarrow RCl + HCl + B$$

- (1) POCl<sub>3</sub> and H<sub>3</sub>PO<sub>3</sub>
- (2) POCl<sub>3</sub> and H<sub>3</sub>PO<sub>4</sub>
- (3) H<sub>3</sub>PO<sub>4</sub> and POCl<sub>3</sub>
- (4) H<sub>3</sub>PO<sub>3</sub> and POCl<sub>3</sub>

Ans. (4)

Mass in grams of copper deposited by passing 88. 9.6487 A current through a voltmeter containing copper sulphate solution for 100 seconds is:

(Given: Molar mass of Cu:  $63 \text{ g mol}^{-1}$ , 1F = 96487 C)

- (1) 3.15 g
- (2) 0.315 g
- (3) 31.5 g
- (4) 0.0315 g

Ans. (2)

**89.** The plot of osmotic pressure  $(\Pi)$  vs concentration (mol L<sup>-1</sup>) for a solution gives a straight line with slope 25.73 L bar mol<sup>-1</sup>. The temperature at which the osmotic pressure measurement is done is:

(Use  $R = 0.083 L bar mol^{-1} K^{-1}$ )

- (1) 37℃
- (2) 310°C
- (3) 25.73°C
- (4) 12.05°C

Ans. (1)

90. Identify the major product C formed in the following reaction sequence:

$$\begin{array}{c} CH_{3} - CH_{2} - CH_{2} - I \xrightarrow{\text{NaCN}} A \\ \xrightarrow{\text{OH}^{-}} B \xrightarrow{\text{NaOH}} C \\ \xrightarrow{\text{Partial hydrolysis}} B \xrightarrow{\text{NaOH}} C \\ \xrightarrow{\text{Major}} C \\ \end{array}$$


- (1) propylamine
- (2) butylamine
- (3) butanamide
- (4)  $\alpha$  bromobutanoic acid

Ans. (1)

- **91.** Identify the **correct** answer.
  - (1) Three resonance structures can be drawn for ozone
  - (2) BF<sub>3</sub> has non-zero dipole moment
  - (3) Dipole moment of NF<sub>3</sub> is greater than that of NH<sub>3</sub>
  - (4) Three canonical forms can be drawn for  $CO_3^{2-}$ ion.

Ans. (4)





eedge

**Statement I**:  $\left[\operatorname{Co}(\operatorname{NH_3})_6\right]^{3^+}$  is a homoleptic complex whereas  $\left[\operatorname{Co}(\operatorname{NH_3})_4\operatorname{Cl_2}\right]^+$  is a heteroleptic complex.

**Statement II**: Complex  $\left[\text{Co(NH}_3)_6\right]^{3+}$  has only one kind of ligands but  $\left[\text{Co(NH}_3)_4\text{Cl}_2\right]^+$  has more than one kind of ligands.

In the light of the above statements, choose the *correct* answer from the options given below.

- (1) Both Statement I and Statement II are true.
- (2) Both Statement I and Statement II are false.
- (3) Statement I is true but Statement II is false.
- (4) Statement I is false but Statement II is true.

## Ans. (1)

## 93. For the given reaction

$$\begin{array}{c|c}
C = CH \xrightarrow{KMnO_a/H^*} P' \\
H & & \\
\end{array}$$
(major product)

'P' is

Ans. (2)

- (1) Ce<sup>4+</sup> and Yb<sup>2+</sup>
- (2) Ce<sup>3+</sup> and Eu<sup>2+</sup>
- (3)  $Gd^{3+}$  and  $Eu^{3+}$
- (4) Pm<sup>3+</sup> and Sm<sup>3+</sup>

#### Ans. (1)

**95.** Consider the following reaction in a sealed vessel at equilibrium with concentrations of

$$N_2 = 3.0 \times 10^{-3} \, M, \, O_2 = 4.2 \times 10^{-3} \, M$$
 and

$$NO = 2.8 \times 10^{-3} M.$$

$$2NO_{(g)} \rightleftharpoons N_{2(g)} + O_{2(g)}$$

If  $0.1 \text{ mol } L^{-1}$  of  $NO_{(g)}$  is taken in a closed vessel, what will be degree of dissociation (a) of  $NO_{(g)}$  at equilibrium?

- (1) 0.00889
- (2) 0.0889
- (3) 0.8889
- (4) 0.717

## Ans. (4)

**96.** A compound X contains 32% of A, 20% of B and remaining percentage of C. Then, the empirical formula of X is:

(Given atomic masses of A = 64; B = 40; C = 32 u)

- (1)  $A_2BC_2$
- (2) ABC<sub>3</sub>
- (3)  $AB_2C_2$
- (4) ABC<sub>4</sub>

## Ans. (2)

**97.** The work done during reversible isothermal expansion of one mole of hydrogen gas at 25°C from pressure of 20 atmosphere to 10 atmosphere is:

(Given  $R = 2.0 \text{ cal } K^{-1} \text{ mol}^{-1}$ )

- (1) 0 calorie
- (2) -413.14 calories
- (3) 413.14 calories
- (4) 100 calories

Ans. (2)







- During the preparation of Mohr's salt solution 98. (Ferrous ammonium sulphate), Which of the following acid is added to prevent hydrolysis of Fe<sup>2+</sup> ion?
  - (1) dilute hydrochloric acid
  - (2) concentrated sulphuric acid
  - (3) dilute nitric acid
  - (4) dilute sulphuric acid

Ans. (4)

The rate of a reaction quadruples when temperature **99**. changes from 27°C to 57°C.


Calculate the energy of activation.

Given  $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$ ,  $\log 4 = 0.6021$ 

- (1) 38.04 kJ/mol
- (2) 380.4 kJ/mol
- (3) 3.80 kJ/mol
- (4) 3804 kJ/mol

ARE YOU NEE

Ans. (1)



$$H_3C$$
 $PBr_3$ 
 $A$ 
 $alc.KOH$ 
 $\Delta$ 
 $B$ 
 $(major)$ 

(1) 
$$A = H_3C$$

$$H_3C$$

$$H_3C$$

$$B = H_3C$$

(3) 
$$A = H_3C$$

$$H_3C$$

$$Br$$

$$B = H_3C$$

(4) 
$$A = H_3C$$

$$H_3C$$

$$B = H_3C$$

$$B = H_3C$$